

N(A)-TERNARY SEMIGROUPS

Ch. Manikya Rao¹, P. Koteswara Rao² and D. Madhusudhana Rao³

¹ Department of Mathematics, Bapatla Arts & Sciences College,
Bapatla, Guntur (Dt), A. P. India. Email : cmrao1998@gmail.com

² Acharya Nagarjuna University, Nagarjuna Nagar, Guntur

³ Department of Mathematics, V. S. R & N. V. R. College, Tenali, A. P. India.

Email : dmrmaths@gmail.com

ABSTRACT

In this paper, the terms, 'A-potent', 'left A-divisor', 'right A-divisor', 'A-divisor' elements, 'N(A)-ternary semigroup' for an ideal A of a ternary semigroup are introduced. If A is an ideal of a ternary semigroup T then it is proved that (1) $A \subseteq N_2(A) \subseteq N_1(A) \subseteq N_0(A)$ (2) $N_0(A) = A_2$, $N_1(A)$ is a semiprime ideal of T containing A, $N_2(A) = A_4$ are equivalent, where $N_0(A)$ = The set of all A-potent elements in T, $N_1(A)$ = The largest ideal contained in $N_0(A)$, $N_2(A)$ = The union of all A-potent ideals. If A is a semipseudo symmetric ideal of a ternary semigroup then it is proved that $N_0(A) = N_1(A) = N_2(A)$. It is also proved that if A is an ideal of a ternary semigroup such that $N_0(A) = A$ then A is a completely semiprime ideal. Further it is proved that if A is an ideal of ternary semigroup T then $R(A)$, the divisor radical of A, is the union of all A-divisor ideals in T. In a N(A)- ternary semigroup it is proved that $R(A) = N_1(A)$. If A is a semipseudo symmetric ideal of a semigroup T then it is proved that S is an N(A)- ternary semigroup iff $R(A) = N_0(A)$. It is also proved that if M is a maximal ideal of a ternary semigroup T containing a pseudo symmetric ideal A then M contains all A-potent elements in T or $T \setminus M$ is singleton which is A-potent.

Mathematical subject classification (2010) : 20M07; 20M11; 20M12.

KEY WORDS: Pseudo symmetric ideal, semipseudo symmetric ideal, prime ideal, semiprime ideal, completely prime ideal, completely semiprime ideal, semisimple element, A-potent element, A-potent ideal, A-divisor, N(A)- ternary semigroup.

1. INTRODUCTION :

The theory of ternary algebraic systems was introduced by LEHMER [12] in 1932, but earlier such structures were studied by KERNER [11] who gave the idea of n -ary algebras. LEHMER investigated certain algebraic systems called triplexes which turn out to be commutative ternary groups. Ternary semigroups are universal algebras with one associative ternary operation. The notion of ternary semigroup was known to BANACH who is credited with an example of a ternary semigroup which cannot reduce to a semigroup. LOS [13] studied some properties of ternary semigroups and proved that every ternary semigroup can be embedded in a semigroup. SIOSON [22] introduced the ideal theory in ternary semigroups. He also introduced the notion of regular ternary semigroups and characterized them by using the properties of quasi-ideals. SANTIAGO developed the theory of ternary semigroups and semiheaps. He studied regular and completely regular ternary semigroups. DIXIT and DEWAN studied quasi-ideals and bi-ideals in ternary semigroups. M. SHABIR and A. KHAN studied prime ideals and prime one-sided ideals in semigroups. SHABIR and BASHIR launched prime ideals in ternary semigroups. In this paper we introduce the notion of $N(A)$ -ternary semigroup and characterize $N(A)$ -ternary semigroups.

2. PRELIMINARIES :

DEFINITION 2.1 : Let T be a non-empty set. Then T is said to be a *ternary semigroup* if there exists a mapping from $T \times T \times T$ to T which maps $(x_1, x_2, x_3) \rightarrow x_1 x_2 x_3$ satisfying the condition : $[x_1 x_2 x_3 \ x_4 x_5] = [x_1 \ x_2 x_3 x_4 \ x_5] = [x_1 x_2 \ x_3 x_4 x_5] \ \forall x_i \in T, 1 \leq i \leq 5$.

DEFINITION 2.2 : A ternary semigroup T is said to be *commutative* provided for all $a, b, c \in T$, we have $abc = bca = cab = bac = cba = acb$.

DEFINITION 2.3 : Let T be a ternary semigroup. A non-empty subset S of T is said to be a *ternary subsemigroup* of T if $abc \in S$ for all $a, b, c \in S$.

NOTE 2.4 : A non-empty subset S of a ternary semigroup T is a ternary subsemigroup if and only if $SSS \subseteq S$.

DEFINITION 2.5 : A non-empty subset A of a ternary semigroup T is said to be *left ternary ideal* or *left ideal* of T if $b, c \in T, a \in A$ implies $bca \in A$.

DEFINITION 2.6 : A nonempty subset of a ternary semigroup T is said to be a *lateral ternary ideal* or simply *lateral ideal* of T if $b, c \in T, a \in A$ implies $bac \in A$.

DEFINITION 2.7 : A nonempty subset A of a ternary semigroup T is a *right ternary ideal* or simply *right ideal* of T if $b, c \in T, a \in A$ implies $abc \in A$

DEFINITION 2.8 : An ideal A of a ternary semigroup T is said to be a *completely prime ideal* of T provided $x, y, z \in T$ and $xyz \in A$ implies either $x \in A$ or $y \in A$ or $z \in A$.

DEFINITION 2.9 : An ideal A of a ternary semigroup T is said to be a *prime ideal* of T provided X,Y,Z are ideals of T and $XYZ \subseteq A \Rightarrow X \subseteq A$ or $Y \subseteq A$ or $Z \subseteq A$.

DEFINITION 2.10 : An ideal A of a ternary semigroup T is said to be a *completely semiprime ideal* provided $x \in T, x^n \in A$ for some odd natural number $n > 1$ implies $x \in A$.

DEFINITION 2.11 : An ideal A of a ternary semigroup T is said to be *semiprime ideal* provided X is an ideal of T and $X^n \subseteq A$ for some odd natural number n implies $X \subseteq A$.

DEFINITION 2.12 : An ideal A of a ternary semigroup T is said to be *pseudo symmetric* provided $x, y, z \in T, xyz \in A$ implies $xsytz \in A$ for all $s, t \in T$.

THEOREM 2.13 : Every completely semiprime ideal A in a ternary semigroup T is a pseudo symmetric ideal.

DEFINITION 2.14 : An ideal A in a ternary semigroup T is said to be *semipseudo symmetric* provided for any odd natural number $n, x \in T, x^n \in A \Rightarrow \langle x \rangle^n \subseteq A$.

NOTATION 2.15 : If A is an ideal of a ternary semigroup T, then we associate the following four types of sets.

A_1 = The intersection of all completely prime ideals of T containing A.

$A_2 = \{x \in T: x^n \in A \text{ for some odd natural numbers } n\}$

A_3 = The intersection of all prime ideals of T containing A.

$A_4 = \{x \in T: \langle x \rangle^n \subseteq A \text{ for some odd natural number } n\}$

THEOREM 2.16: If A is an ideal of a ternary semigroup T, then $A \subseteq A_4 \subseteq A_3 \subseteq A_2 \subseteq A_1$.

THEOREM 2.17 : Let A be a semipseudo symmetric ideal of a ternary semigroup T. Then the following are equivalent.

- 1) A_1 = The intersection of all completely prime ideals of T containing A.
- 2) A_1^1 = The intersection of all minimal completely prime ideals of T containing A.
- 3) A_1^{11} = The minimal completely semiprime ideal of T relative to containing A.
- 4) A_2 = $\{x \in T : x^n \in A \text{ for some odd natural number } n\}$
- 5) A_3 = The intersection of all prime ideals of T containing A.
- 6) A_3^1 = The intersection of all minimal prime ideals of T containing A.
- 7) A_3^{11} = The minimal semiprime ideal of T relative to containing A.
- 8) A_4 = $\{x \in T : \langle x \rangle^n \subseteq A \text{ for some odd natural number } n\}$

3. N(A)-TERNARY SEMIGROUPS :

DEFINITION 3.1 : Let A be an ideal in a Ternary semigroup T. An element $x \in T$ is said to be *A-potent* there exists a odd natural number n such that $x^n \in A$.

DEFINITION 3.2 : Let A be an ideal in a Ternary semigroup T. An ideal B of T is said to be *A-potent ideal* provided there exists a odd natural number n such that $B^n \subseteq A$.

NOTE 3.3 : If A is an ideal of a ternary semigroup T, then every element of A is a A-potent element of T and A itself an A-potent ideal of T.

DEFINITION 3.4 : Let A be an ideal of a ternary semigroup T. An A-potent element x is said to be a *nontrivial A-potent* element of T if $x \notin A$.

NOTATION 3.5 : $N_0(A)$ = The set of all A-potent elements in T.

$N_1(A)$ = The largest ideal contained in $N_0(A)$.

$N_2(A)$ = The union of all A-potent ideals.

THEOREM 3.6 : If A is an ideal of a ternary semigroup T, then $A \subseteq N_2(A) \subseteq N_1(A) \subseteq N_0(A)$.

Proof : Since A is itself an A-potent ideal, and $N_2(A)$ is the union of all A-potent ideals. Therefore $A \subseteq N_2(A)$. Let $x \in N_2(A) \Rightarrow x$ belongs to at least one A-potent ideal $\Rightarrow x$ is an A-potent element. Hence $x \in N_0(A)$. Therefore $N_2(A) \subseteq N_0(A)$. Clearly $N_2(A)$ is an ideal of T. Since $N_1(A)$ is the largest ideal contained in $N_0(A)$, we have $N_2(A) \subseteq N_1(A) \subseteq N_0(A)$. Hence $A \subseteq N_2(A) \subseteq N_1(A) \subseteq N_0(A)$.

THEOREM 3.7 : If A is an ideal in a ternary semigroup T , then the following are true.

1. $N_0(A) = A_2$.
2. $N_1(A)$ is a semiprime ideal of S containing A .
3. $N_2(A) = A_4$.

Proof : (1) $N_0(A)$ = The set of all A -potent elements

$$= \{ x \in S : x^n \in A \text{ for some natural number } n \} = A_2.$$

(2) Suppose that $\langle x \rangle^2 \subseteq N_1(A)$. Suppose if possible $x \notin N_1(A)$. $N_1(A)$, $\langle x \rangle$ are the ideals implies $N_1(A) \cup \langle x \rangle$ is an ideal. Since $N_1(A)$ is the largest ideal in $N_0(A)$,

We have $N_1(A) \cup \langle x \rangle \not\subseteq N_0(A) \Rightarrow \langle x \rangle \not\subseteq N_0(A)$. Hence there exists an element y such that

$y \in \langle x \rangle \setminus N_0(A)$. Now $y^3 \in \langle x \rangle^3 \subseteq N_1(A) \subseteq N_0(A) \Rightarrow y^3 \in N_0(A) \Rightarrow (y^3)^n \in A$ for some odd natural number $n \Rightarrow y^{3n} \in A \Rightarrow y \in N_0(A)$. It is a contradiction. Therefore $x \in N_1(A)$. Hence $N_1(A)$ is a semiprime ideal of S containing A .

(3) Let $x \in N_2(A)$. Then there exists an A -potent ideal B such that $x \in B$. B is A -potent ideal implies there exists an odd natural number $n \in \mathbb{N}$ such that $B^n \subseteq A \Rightarrow \langle x \rangle^n \subseteq B^n \subseteq A$ for some $n \in \mathbb{N} \Rightarrow x \in A_4$. Therefore $N_2(A) \subseteq A_4$. Let $x \in A_4 \Rightarrow \langle x \rangle^n \subseteq A$ for some $n \in \mathbb{N}$. So $\langle x \rangle$ is an A -potent ideal in S and hence $\langle x \rangle \subseteq N_2(A) \Rightarrow x \in N_2(A)$. Therefore $A_4 \subseteq N_2(A)$. Hence $N_2(A) = A_4$.

NOTE 3.8 : It is natural to ask whether $N_1(A) = A_3$. This is not true.

EXAMPLE 3.9 : In the free semigroup S over the alphabet a, b, c . For the ideal $A = Sa^3S$, $N_0(A) = \{a\} \cup S^1a^3S^1$ and $N_1(A) = \{a^3\} \cup Sa^3S = S^1a^3S^1$. But Sa^3S is a prime ideal, let A, B, C are three ideals of S such that $ABC \subseteq Sa^3S$, implies all words containing $a^3 \in A$ or all words containing $a^3 \in B$ or all words containing $a^3 \in C \Rightarrow A \subseteq Sa^3S$ or $B \subseteq Sa^3S$ or $C \subseteq Sa^3S$. Therefore Sa^3S is a prime ideal. We have $A_3 = Sa^3S$, so $N_1(A) \neq A_3$. Therefore we can remark that the inclusions in $A_3 \subseteq N_1(A) \subseteq N_0(A) = A_2$ may be proper in an arbitrary ternary semigroup.

THEOREM 3.10 : If A is a semipseudo symmetric ideal in a ternary semigroup T , then $N_0(A) = N_1(A) = N_2(A)$.

Proof : Suppose A is a semi pseudo symmetric ideal in a ternary semigroup T . By theorem 3.7, $N_0(A) = A_2$ and $N_2(A) = A_4$. Also by theorem 2.17, we have $A_2 = A_4$. Hence

$N_0(A) = N_2(A)$. By the theorem 3.6, $A \subseteq N_2(A) \subseteq N_1(A) \subseteq N_0(A)$. We have $N_2(A) \subseteq N_1(A)$. Now let $x \in N_1(A) \Rightarrow x \in N_0(A) \Rightarrow x \in N_2(A)$. Therefore $N_1(A) \subseteq N_2(A)$. Hence $N_1(A) = N_2(A)$. Therefore $N_0(A) = N_1(A) = N_2(A)$.

THEOREM 3.11 : For any semipseudo symmetric ideal A in a ternary semigroup T , a nontrivial A -potent element $x(x \notin A)$ cannot be semisimple.

Proof : Since x is a nontrivial A -potent element, there exists a odd natural number n such that $x^n \in A$. Since A is semipseudo symmetric ideal, we have $\langle x \rangle^n \subseteq A$. If x is semisimple, then $\langle x \rangle = \langle x \rangle^3$ and hence $\langle x \rangle = \langle x \rangle^n \subseteq A$, this is a contradiction. Thus x is not semisimple.

THEOREM 3.12 : If A is an ideal in a ternary semigroup T , such that $N_0(A) = A$, then A is a completely semiprime ideal and A is a pseudo symmetric ideal.

Proof: Let $x \in T$ and $x^3 \in A$. Since $N_0(A) = A$, $x^3 \in N_0(A)$. Thus there exists a odd natural number n such that $(x^3)^n \in A \Rightarrow x^{3n} \in A \Rightarrow x \in N_0(A) = A$. Therefore A is a completely semiprime ideal. By theorem 2.12, A is pseudo symmetric ideal. Hence A is completely semiprime and pseudo symmetric ideal.

THEOREM 3.13 : If A is a semipseudo symmetric ideal of a ternary semisimple semigroup the $A = N_0(A)$.

Proof : Clearly $A \subseteq N_0(A)$. Let $a \in N_0(A)$. If $a \notin A$ then a is a nontrivial A -potent element. By theorem 3.11, a can not be semisimple. It is a contradiction. Therefore $a \in A$ and hence $N_0(A) \subseteq A$. Thus $N_0(A) = A$.

DEFINITION 3.114 : Let A be an ideal in a ternary semigroup T . An element $x \in T$ is said to be a *left A-divisor* provided there exists an element $y, z \in T \setminus A$ such that $xyz \in A$.

DEFINITION 3.15 : Let A be an ideal in a ternary semigroup T . An element $x \in T$ is said to be a *right A-divisor* provided there exists an element $y, z \in T \setminus A$ such that $yzx \in A$.

DEFINITION 3.16 : Let A be an ideal in a ternary semigroup T . An element $x \in T$ is said to be an *A-divisor* if x is both a left A -divisor and a right A -divisor element.

DEFINITION 3.17 : Let A be an ideal in a ternary semigroup T . An ideal B in T is said to be a *left A-divisor ideal* provided every element of B is a left A -divisor element.

DEFINITION 3.18 : Let A be an ideal in a ternary semigroup T . An ideal B in T is said to be a *right A-divisor ideal* provided every element of B is a right A -divisor element.

DEFINITION 3.19 : Let A be an ideal in a ternary semigroup T . An ideal B in T is said to be an *A-divisor ideal* provided if it is both a left A -divisor ideal and a right A -divisor ideal of a semigroup T .

NOTATION 3.20 : $R_l(A)$ = The union of all left A -divisor ideals in T .

$R_r(A)$ = The union of all right A -divisor ideals in T .

$R(A) = R_l(A) \cap R_r(A)$. We call $R(A)$, the divisor radical of T .

THEOREM 3.21: If A is any ideal of a ternary semigroup T , then $N_1(A) \subseteq R(A)$.

Proof : Let $x \in N_1(A)$. Since $N_1(A) \subseteq N_0(A)$, we have $x \in N_0(A) \Rightarrow x^n \in A$ for some odd natural number n . Let n be the least odd natural number such that $x^n \in A$. If $n = 1$ then $x \in A$ and hence $x \in R(A)$.

If $n > 1$, then $x^n = x^{n-2} \cdot x \in A$, where $x^{n-2} \in T \setminus A$. Hence x is an A -divisor element.

Thus $x \in R(A)$. Therefore $N_1(A) \subseteq R(A)$.

THEOREM 3.22 : If A is an ideal in a ternary semigroup T , then $R(A)$ is the union of all A -divisor ideals in T .

Proof : Suppose A is an ideal in a ternary semigroup T . Let B be an A -divisor ideal in T . Then B is both a left A -divisor and a right A -divisor ideal in T . Thus $B \subseteq R_l(A)$ and $B \subseteq R_r(A) \Rightarrow B \subseteq R_l(A) \cap R_r(A) = R(A) \Rightarrow B \subseteq R(A)$. Therefore $R(A)$ contains the union of all A -divisor ideals in T . Let $x \in R(A)$. Then $x \in R_l(A) \cap R_r(A)$. So $\langle x \rangle \subseteq R_l(A) \cap R_r(A)$. Hence $\langle x \rangle$ is an A -divisor ideal. So $R(A)$ is contained in the union of all divisor ideals in T . Thus $R(A)$ is the union of all divisor ideals of T .

COROLLARY 3.23 : If A is a pseudo symmetric ideal in a semigroup T , then $R(A)$ is the set of all A -divisor elements in T .

Proof : Suppose A is a pseudo symmetric ideal in T . Let x be an A -divisor element in T . Then $xyz \in A$, where $y, z \in S \setminus A$. $xyz \in A$, A is pseudo symmetric $\Rightarrow \langle x \rangle \langle y \rangle \langle z \rangle \subseteq A \Rightarrow \langle x \rangle$ is an A -divisor ideal $\Rightarrow \langle x \rangle \subseteq R(A) \Rightarrow x \in R(A)$. Hence $R(A)$ is the set of all A -divisor elements in T .

DEFINITION 3.24 : Let A be an ideal in a ternary semigroup T . T is said to be a $N(A)$ -ternary semigroup provided every A -divisor is A -potent.

NOTATION 3.25 : Let T be a semigroup with zero. If $A=\{0\}$, then we write R for $R(A)$ and N for $N_0(A)$ and N - ternary semigroup for $N(A)$ -ternary semigroup.

THEOREM 3.26 : If T is an $N(A)$ -ternary semigroup, then $R(A) = N_1(A)$.

Proof : Suppose S is an $N(A)$ -ternary semigroup. By theorem 3.21, $N_1(A) \subseteq R(A)$.

Let $x \in R(A) \Rightarrow x$ is an A -divisor $\Rightarrow x$ is an A -potent $\Rightarrow x \in N_1(A)$. $\therefore R(A) \subseteq N_1(A)$.

Hence $N_1(A) = R(A)$.

THEOREM 3.27 : Let A be a semipseudo symmetric ideal in a ternary semigroup T . Then T is an $N(A)$ -ternary semigroup iff $R(A) = N_0(A)$.

Proof : Since A is a semipseudo symmetric ideal, by theorem 3.10, $N_0(A) = N_1(A) = N_2(A)$. If T an $N(A)$ -ternary semigroup, then by theorem 3.26, $R(A) = N_1(A)$. Hence $R(A) = N_0(A)$. Conversely suppose that $R(A) = N_0(A)$. Then clearly every A -divisor element is an A -potent element. Hence T is an $N(A)$ -ternary semigroup.

COROLLARY 3.28 : Let A be a pseudo symmetric ideal in a ternary semigroup T . Then T is an $N(A)$ -ternary semigroup if and only if $R(A) = N_0(A)$.

Proof : Since every pseudo symmetric ideal is a semipseudo symmetric ideal, by theorem 3.27, $R(A) = N_0(A)$.

COROLLARY 3.29 : Let T be a semigroup with 0 and $\langle 0 \rangle$ is a pseudo symmetric ideal. Then $R = N$ iff T is an N -ternary semigroup.

Proof : The proof follows from the theorem 3.24.

THEOREM 3.30: If M is a maximal ideal in a ternary semigroup T containing a pseudo symmetric ideal A , then M contains all A -potent elements in T or $T \setminus M$ is singleton which is A -potent.

Proof : Suppose M does not contain all A -potent elements.

Let $a \in T \setminus M$ be any A -potent element and b be any element in $T \setminus M$.

Since M is a maximal ideal, $M \cup \langle a \rangle = M \cup \langle b \rangle \Rightarrow \langle a \rangle = \langle b \rangle$.

Since $b \notin M$, we have $b \in \langle a \rangle$. Let n be the least positive odd integer such that $a^n \in A$. Since A is a pseudo symmetric ideal then A is a semipseudo symmetric ideal and hence $\langle a \rangle^n \subseteq A$. Therefore $b^n \in A$ and hence b is an A -potent element.

Thus every element in $T \setminus M$ is A -potent.

Similarly we can show that if m is the least positive odd integer such that $b^m \in A$, then $a^m \in A$. Therefore there exists an odd natural number p such that $a^p \in A$ and $a^{p-2} \notin A$ for all $a \in T \setminus M$. Let $a, b, c \in T \setminus M$.

Since M is maximal ideal, we have $\langle a \rangle = \langle b \rangle = \langle c \rangle$.

So $b, c \in \langle a \rangle \Rightarrow b = sat, c = uav$. So $a \in \langle b \rangle$ and hence $a = sbt$ for some $s, t \in T^1$.

Now since A is a pseudo symmetric ideal,

we have $(abc)^{p-2} = (abc)^{p-3}abc = (abc)^{p-3}a(sat)(uav) \in A \Rightarrow abc \in M$.

If $b \neq a$ then $s, t \in T$. If $s, t \in M$ then $sat \in M \Rightarrow b \in M$. Which is not true.

In both the cases we have a contradiction. Hence $a = b$. Similarly we show that $c = a$.

COROLLARY 3.31 : If M is a nontrivial maximal ideal in a ternary semigroup T containing a pseudo symmetric ideal A . Then $N_0(A) \subseteq M$.

Proof : Suppose in $N_0(A) \not\subseteq M$. Then by above theorem 3.30, M is trivial ideal. It is a contradiction. Therefore $N_0(A) \subseteq M$.

COROLLARY 3.32 : If M is a maximal ideal in a semisimple ternary semigroup T containing a semipseudo symmetric ideal A . Then $N_0(A) \subseteq M$.

Proof : By theorem 3.13, A is pseudo symmetric ideal. If $a \in T \setminus M$ is A -potent, then a cannot be semisimple. It is a contradiction. Therefore $N_0(A) \subseteq M$.

REFERENCES:

- [1] ANJANEYULU A., *Primary ideals in semigroups* – Semigroup Form, Vol.20 (1980) 129-144.
- [2] ANJANEYULU A., *On primary semigroups* – Czechoslovak Mathematical Journal, 30(105) 1980, Praha.
- [3] ANJANEYULU A., *Semigroups in which prime ideals are maximal* – Semigroup Form, Vol.22 (1981) 151-158.
- [4] ANJANEYULU A., *Structure and ideal theory of duo semigroups* – Semigroup Form Vol.22 (1981) 257-276.
- [5] ANJANEYULU A., *Structure and ideal theory of semigroups* – Thesis, ANU (1980).
- [6] DIXIT. V. N and DEWAN. S., *A note on quasi and bi-ideals in ternary semigroups*, Int. J. Math. Sci. 18 (1995), 501 – 508.
- [7] DIXIT. V. N and DEWAN. S., *Minimal quasi-ideals in ternary semigroups*, Indian J. Pure Appl. Math. 28 (1997), 255 – 632.
- [8] HEWITT. E and JZUCKERMAN. H.S., *Ternary Operations and semigroups*- Semigroup Proc. Sympos, Wayne State Univ., Detroit, 1968, 55-83.
- [9] IAMPAN. A., *On ordered ideal extensions of ordered ternary semigroup*- Lobachevskii Journal of Mathematics, 31(1)(2010), 13-17.
- [10] IAMPAN. A., *Lateral ideals of ternary semigroups* – Yxpaihckkhii Mate. Math. Torn 4 (2007) no.4, 525-534.
- [11] KERNER. R., *Ternary Algebraic Structures and their applications*-Univ, P & M, Qurie pre print, Paris (2000), Arixiv Math.
- [12] LEHMER. D. H., *A ternary analogue of abelian groups*, Amer. J. Math., 59(1932), 329-338.
- [13] LYAPIN. E. S., *Realization of ternary semigroups* – Russian Modern Algebra, Leningrad University, Leningrad, 1981, pp 43-48.
- [14] LYAPIN. E. S., *Realization of ternary semigroups* – Russian Modern Algebra, Leningrad University, Leningrad, 1981, pp 43-48.
- [15] MEHDI DEGHANIAN and MOHAMMAD SADEGH MODARRES – *Ternary γ -homomorphisms and ternary γ -derivations on ternary semigroups* – Journal of Inequalities and applications 2012 : 34.
- [16] MOHAMMAD SHABIR and MEHAR BANO- *Prime Bi-ideals in Ternary semigroups, Quasi groups and Related systems (16)* (2008) 239-256.
- [17] SARALA. Y, ANJANEYULU. A and MADHUSUDHANA RAO.D., *Ternary semigroups*, - International Journal of Mathematical Sciences, Technology and Humanities 76 (2013) 848-859.
- [18] SARALA. Y, ANJANEYULU. A and MADHUSUDHANA RAO.D., *ideals in ternary semigroups* - International eJournal of Mathematics and Engineering 203 (2013) 1950-1968.
- [19] SARALA. Y, ANJANEYULU. A and MADHUSUDHANA RAO. D., *Prime radicals in ternary semigroups* - international organization of scientific research journal of Mathematics ISSN: 2278-5728. Volume 4, Issue 5 (Jan.-Feb. 2013), pp 43-53.
- [20] SARALA. Y, ANJANEYULU. A and MADHUSUDHANA RAO. D., *pseudo symmetric ideals in ternary semigroups* - International Referred Journal of Engineering and Science ISSN (Online) 2319-183X, (print) 2319-1821 Volume 1, Issue 4 (December 2012), pp. 33-43.

- [21] SARALA. Y, ANJANEYULU. A and MADHUSUDHANA RAO. D., *Pseudo Integral ternary semigroups* –Asian Journal of Current Engineering and Maths 2:1 Jan-Feb (2013) 40-43.
- [22] SIOSON. F. M., *Ideal theory in ternary semigroups*, Math. Japon.10(1965), 63-84.

Author's Personal Copy